



BRIEF REPORT

Epidemiological and molecular characteristics of *Chlamydia psittaci* from 8 human cases of psittacosis and 4 related birds in Argentina



María E. Cadario^{a,*},[◇], María C. Frutos^{b,◇}, Maite B. Arias^a, Javier A. Origlia^c, Vanina Zelaya^d, María J. Madariaga^e, Claudia S. Lara^a, Viviana Ré^b, Cecilia G. Cuffini^b

^a INEI-ANLIS «Dr. Carlos G. Malbrán», Ciudad Autónoma de Buenos Aires, Argentina

^b Instituto de Virología «Dr. J.M. Vanella», Facultad de Ciencias Médicas, Universidad Nacional de Córdoba, Córdoba, Argentina

^c Cátedra de Patología de Aves y Pilíferos, Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata, Argentina

^d UOCCB-ANLIS «Dr. Carlos G. Malbrán», Ciudad Autónoma de Buenos Aires, Argentina

^e Instituto de Zoonosis Luis Pasteur, Departamento de Diagnóstico y Producción de Biológicos, Ciudad Autónoma de Buenos Aires, Argentina

Received 1 August 2016; accepted 10 April 2017

Available online 19 July 2017

KEYWORDS

Chlamydia psittaci;
Genotype A;
Psittacosis

Abstract In Argentina, the epidemiological and molecular characteristics of *Chlamydia psittaci* infections are still not sufficiently known. A total of 846 respiratory and 10 ocular samples from patients with suspected human psittacosis were tested for *C. psittaci* from January 2010 to March 2015. Four samples of birds related to these patients were also studied. Forty-eight samples were positive for *C. psittaci* by a nested PCR. The molecular characterization of twelve *C. psittaci* PCR-positive samples received in the National Reference Laboratory INEI-ANLIS «Dr. Carlos G. Malbrán», Buenos Aires, Argentina was performed. Eight positive samples from humans and four from birds were genotyped by *ompA* gene sequencing. *C. psittaci* genotype A was found in all human samples and in the related birds. This report contributes to our increasing knowledge of the epidemiological and molecular characteristics of *C. psittaci* to conduct effective surveillance of its zoonotic infections.

© 2017 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: ecadario@anlis.gov.ar (M.E. Cadario).

◇ These authors contributed equally to this article.

PALABRAS CLAVE
Chlamydia psittaci;
 Genotipo A;
 Psittacosis

Características epidemiológicas y moleculares de *Chlamydia psittaci* provenientes de 8 casos humanos de psittacosis y de 4 aves relacionadas en la Argentina

Resumen En la Argentina, aún no se conocen suficientemente las características epidemiológicas y moleculares de las infecciones por *Chlamydia psittaci*. Entre enero del 2010 y marzo del 2015 se estudiaron 846 muestras respiratorias y 10 oculares de pacientes con sospecha de psittacosis para la búsqueda de *C. psittaci*. También se estudiaron 4 muestras de aves relacionadas con estos pacientes. De ese total, 48 muestras fueron positivas para *C. psittaci* mediante una reacción en cadena de la polimerasa (PCR) anidada. Posteriormente, se realizó en el INEI-ANLIS «Dr. Carlos G. Malbrán» la caracterización molecular de 12 muestras positivas para *C. psittaci*, 8 de humanos y 4 de aves, que fueron genotipificadas por secuenciación del gen *ompA*. *C. psittaci* genotipo A se encontró en todas esas muestras. Este informe contribuye a mejorar nuestro conocimiento de las características epidemiológicas y moleculares de *C. psittaci* para lograr una vigilancia efectiva de la zoonosis que produce.

© 2017 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Psittacosis is a zoonosis caused by *Chlamydia psittaci*, an obligate intracellular bacterium belonging to the *Chlamydiaceae* family and its single genus, *Chlamydia*¹³.

Human infections vary from unapparent to severe systemic diseases: influenza – like illnesses, severe atypical pneumonia, endocarditis, myocarditis, meningitis, conjunctivitis, and others. The disease is rarely fatal in properly treated patients. Therefore, awareness of an early diagnosis is important. Pulmonary involvement is common and information of recent exposure to birds is frequently omitted during medical consultation.

Confirmation by molecular methods and genotyping of *C. psittaci* infection using human respiratory samples and animal samples is a matter of both diagnostic and epidemiological relevance⁶.

C. psittaci is divided into at least 9 genotypes (A–F, E/B, M56 and WC). Sequence analysis of the *ompA* gene which encodes the major outer membrane protein (MOMP) gene, is one way to identify all known and eventual new genotypes³.

Genotypes A–F and E/B are associated with birds and genotypes WC and M56 have been found in cattle and muskrats, respectively³. Genotype A is associated with Psittaciformes, B with pigeons, C with ducks and geese¹¹, D with turkeys, E with pigeons and ratites, F with psittacine birds and turkeys⁵, and genotype E/B is mainly associated with ducks, turkeys and pigeons^{3,7}.

The aim of this study was the molecular characterization of *C. psittaci* polymerase chain reaction-positive (PCR-positive) samples received in the National Reference Laboratory INEI-ANLIS "Dr. Carlos G. Malbrán", Buenos Aires, Argentina.

Eight hundred and fifty-six samples from patients of different cities of Argentina, in whom psittacosis was suspected based on clinical presentation and/or history of exposure with infected birds, were analyzed between January 2010 and March 2015. Eight hundred and forty-six were respiratory samples (nasal and pharyngeal swabs, nasopharyngeal aspirates, tracheal aspirates and bronchioalveolar lavages) and 10 were ocular swabs. Cloacal and ocular

swabs from 10 related birds were collected and stored until processed. Dacron swabs were placed in a 2 ml sucrose-phosphate-glutamate or UTM (Copan Italia, Brescia, Italy), and stored at 4 °C. Two hundred microliters of this suspension were subjected to DNA extraction using the DNeasy Blood & Tissue Kit (Qiagen, Valencia, CA, and USA) according to the manufacturer's instructions. DNA was eluted into 200 µl of Qiagen elution buffer and stored at -20 °C until tested.

A nested PCR protocol was used to amplify a portion of the 16S rRNA gene as previously described by Messmer et al.¹⁰ Briefly, 5 µl of DNA extract was used in the first step to amplify a 436 bp fragment common to the *Chlamydia* genus. In a second PCR step, 5 µl of the PCR product obtained in the previous PCR reaction was used to amplify a 127 bp specific product of *C. psittaci*.

Genotyping of *C. psittaci* was attempted in all positive samples in the 16S rRNA gene PCR. Amplification by nested PCR followed by sequencing of the variable domains III and IV of the *ompA* gene of *C. psittaci* was performed as described by Sachse and Hotzel.¹² Briefly, 5 µl of DNA extract were used in the first step to amplify a 576 bp fragment of the *ompA* gene common to the *Chlamydia* genus. In a second PCR step, 5 µl of the PCR product obtained in the previous PCR reaction were used to amplify a 389 bp specific product of *C. psittaci*.

Nested PCR products were purified by gel electrophoresis using the AccuPrep® PCR Purification Kit (USA Bioneer Inc.) and subjected to a direct nucleotide sequencing reaction in both directions using the internal (second-round) PCR primers by Macrogen, Inc. (Seoul, Korea). The sequences obtained from fragments of the *ompA* gene were edited and prepared with BioEdit V 7.0.9⁴ and subsequently aligned with Clustal X 2.12⁸, along with the sequences downloaded from GenBank. Relatedness of newly characterized sequences was assessed by analysis using the 2.2.19 Basic Local Alignment Search Tool. The choice of the genomic region was based on the fact that it is widely associated with the genetic divergence of *C. psittaci* and it identifies all known genotypes³.

The dendrogram was constructed using the Tree Explorer module of the MEGA software version 4¹⁴ with the neighbor-joining method and the *p*-distance parameter. The branch support was evaluated by nonparametric bootstrapping with 1000 pseudo-replicas.

Among the respiratory and ocular swabs samples of the 856 patients studied, 48 (5.6%) tested positive for *C. psittaci* by nested-PCR. Additionally, 6 samples from related birds tested positive.

Among these 54 positive samples for *C. psittaci*, only 12 produced enough PCR products in the *ompA* nested PCR to allow sequencing and genotype determination¹³ (8 from humans and 4 from birds). The mean age of the infected patients was 33.5 years (*r*=5–62 years); 62.5% (5/8) were young workers aged 23–45 years. Male/female ratio was 1:1.

The analysis of the medical records showed that community-acquired pneumonia (CAP) was the most frequent clinical presentation (50%; 4/8), followed by acute respiratory tract infection (25%; 2/8). Whooping cough was suspected in one case (12.5%; 1/8). One patient (12.5%; 1/8) with conjunctivitis tested positive for *C. psittaci*.

Five patients reported contact with birds, especially parrots (*n*=4). Contact with a domestic or companion bird was not reported in the remaining three *C. psittaci* positive patients.

Clinical presentation and sources of the eight *C. psittaci*-positive samples are shown in Table 1.

Genetic diversity and associations among the detected positive samples of *C. psittaci* were determined by the sequencing and genetic analysis of the composition of the *ompA* gene. The sequences obtained in this study were deposited in GenBank under accession numbers KU357040–KU357046; KU357049; KU365350–KU365353.

In the present study all *C. psittaci*-positive samples (from humans and birds) were recognized as belonging to genotype A by phylogenetic analysis (Fig. 1). However, in Córdoba, a central region of Argentina, genotype A was detected in only one case (12.5%, 1/8), being WC (75%, 6/8) the genotype most frequently found in patients with suspected human psittacosis¹ and in captive birds². Probably the differences could be due to different clinical conditions of the patients in both studies or/and to geographical sources. This differential genotypic profile is an interesting finding that should be investigated in the future. *C. psittaci* genotype A is more often found among psittacine birds such as parrots and cockatiels. However, the most prevalent *C. psittaci* genotype in human infections is currently unknown⁶.

The analysis of the medical records showed that CAP was the most frequent clinical presentation. In most studies on CAP, psittacosis was diagnosed only with serological tests. PCR leads to a quick diagnosis and lacks broad genus cross-reactivity. Furthermore, PCR products can be used to genotype *C. psittaci*, which is relevant for public health notification, source detection and control.

Table 1 Characteristics of the *C. psittaci*-positive cases

Patients/birds	Year	City	Sex	Age (years)	Clinical diagnosis	Sample	Contact with birds
67-10	2010	San Miguel	F	62	Community-acquired pneumonia	Tracheal aspirate	Chicken
325-10	2010	Buenos Aires	M	45	Pancreatitis/Community-acquired pneumonia	BAL ^a	Unknown
38-11	2011	Santa Rosa	F	38	Community-acquired pneumonia	N&PS ^a	Parrot
Parrot 38/11A 38-13 ^c	2011 2013	Santa Rosa San Antonio Oeste	NA ^b M	NA ^b 48	NA ^b Acute respiratory tract infection	NA ^b N&PS ^a	NA ^b Illegal parrot trafficking
Parrot 35-13A	2013	San Antonio Oeste	NA ^b	NA ^b	NA ^b	NA ^b	NA ^b
Parrot 38-13A	2013	San Antonio Oeste	NA ^b	NA ^b	NA ^b	NA ^b	NA ^b
Parrot135-13A	2013	San Antonio Oeste	NA ^b	NA ^b	NA ^b	NA ^b	NA ^b
619-13	2013	Trenque Lauquen	F	26	Conjunctivitis	Ocular swab	Pigeons
357-13	2013	Resistencia	M	5	Whooping cough suspected	NPA ^d	Unknown
43-14	2014	Neuquén	F	43	Acute respiratory tract infection	N&PS ^a	Parrot
83-15	2015	Carmen de Areco	M	29	Community-acquired pneumonia	Tracheal aspirate	Unknown

* BAL, bronchioalveolar lavage.

^a N&PS, nasal and pharyngeal swab.

^b NA, not applicable.

^c Index case of an outbreak.

^d NPA, nasopharyngeal aspirate.




Figure 1 Neighbor-joining dendrogram based on comparison of 290 bp fragment of the *ompA* gene in *Chlamydia*. Numbers above branches are bootstrap values as a percentage of 1000 pseudo-replicates and only bootstrap values >60% are shown. *Chlamydia caviae* GPIC was used as an out-group. Scale bar shows the percentage sequence diversity.

The use of nucleic acid amplification directly from clinical samples and direct genotyping with phylogenetic analysis in the *ompA* gene could improve the surveillance and molecular epidemiology of this disease. In this sense, the classical method applying two PCR techniques plus sequencing was successfully used and it should be implemented in regional laboratories for an improved diagnosis of this infectious disease.

In this study, a nested PCR and direct sequencing were used to detect and genetically characterize *C. psittaci*. For the phylogenetic analysis of *C. psittaci*, we used partial *ompA* gene amplification. Our results only report the circulation of genotype A associated with respiratory symptoms and conjunctivitis in different areas of our country, at least in our experience with these eight human cases. Psittaciformes especially parakeets and parrots, and individuals in contact with them or with racing pigeons, were the most frequently infected.

C. psittaci is rarely suspected in ocular infections, but using molecular techniques, as we did, may be more commonly found⁹. In the present study we showed a case of conjunctivitis due to *C. psittaci* in a 26 year-old woman not responding to the current treatment and living in contact with wild birds.

Several limitations of our study need to be considered. Isolation in cell cultures was the preferred technique to confirm and corroborate the results obtained; however, in the case of *C. psittaci*, this procedure is unadvisable because of the biological risks. Moreover, the phylogenetic analysis showed two sequences corresponding to birds (KU365353 and KU365349) in separated clusters. As can be seen in Figure 1 they have a low degree of homology with others obtained in the same city (74% similarity, one with an avian strain and the other with a human strain) (data not shown).

Further studies are needed to confirm the genotype and even the species.

Effective public health surveillance for new variants and modes of transmission will be helpful to understanding the evolution and epidemiology of these bacteria. Therefore, new information about sequence variation of subtypes will be needed.

Ethical responsibilities

Protection of human and animal subjects. The authors state that for this investigation no experiments have been performed on humans or animals.

Confidentiality of data. The authors state that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors state that in this article there are no patient data.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This study was funded by PICTO-ANLIS 2011-0180, Agencia Nacional de Promoción Científica y Tecnológica del Ministerio de Ciencia, Tecnología e Innovación Productiva de la República Argentina. MSc María Estela Cadario is a staff member of the Instituto Nacional de Enfermedades

Infecciosas-ANLIS "Dr Carlos G Malbrán", Buenos Aires, Argentina.

References

1. Frutos MC, Monetti M, Kiguen X, Venezuela F, Ré V, Cuffini C. Genotyping of *C. psittaci* in central area of Argentina. *Diagn Microbiol Infect Dis.* 2012;74:320–2.
2. Frutos MC, Monetti MS, Gallo Vaulet L, Cadario ME, Rodríguez Fermepin M, Ré VE, Cuffini CG. Genetic diversity of *Chlamydia* among captive birds from central Argentina. *Avian Pathol.* 2015;44:50–6.
3. Geens T, Desplanques A, Van Loock M, Bonner BM, Kaleta EF, Magnino S, Andersen AA, Everett KD. Sequencing of *Chlamydia psittaci* *ompA* gene reveals a new genotype, E/B, and need for a rapid discriminatory genotyping method. *J Clin Microbiol.* 2005;43:2456–61.
4. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser.* 1999;41:95–8.
5. Harkinezhad T, Geens T, Vanrompay D. *Chlamydophila psittaci* infections in birds: a review with emphasis on zoonotic consequences. *Vet Microbiol.* 2009;135:68–77.
6. Heddema E, van Hannen E, Duim B, Vandebroucke-Grauls CMJE, Pannekoek Y. Genotyping of *Chlamydophila psittaci* in human samples. *Emerg Infect Dis.* 2006;12:1989–90.
7. Knittler MR, Sachse K. *Chlamydia psittaci*: update on an underestimated zoonotic agent. *Pathog Dis.* 2015;73:1–15.
8. Larkin MA, Blackshields G, Brown NP, Chenna R, McGgettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Clustal W, Clustal. X version 2.0. *Bioinformatics.* 2007;23:2947–8.
9. Lietman T, Brooks D, Moncada J, Schachter J, Dawson C, Dean D. Chronic follicular conjunctivitis associated with *Chlamydia psittaci* or *Chlamydia pneumoniae*. *Clin Infect Dis.* 1998;26:1335–40.
10. Messmer TO, Skelton SK, Moroney JF, Daugherty H, Fields BS. Application of a nested, multiplex PCR to psittacosis outbreaks. *J Clin Microbiol.* 1997;35:2043–6.
11. Pannekoek Y, Dickx V, Beeckman DS, Jolley KA, Keijzers WC, Vretou E, Maiden MC, Vanrompay D, Van der Ende A. Multilocus sequence typing of *Chlamydia* reveals an association between *Chlamydia psittaci* genotypes and host species. *PLoS One.* 2010;5:e14179.
12. Sachse K, Hotzel H. Detection and differentiation of *Chlamydiae* by nested PCR. *Methods Mol Biol.* 2003;216:123–36.
13. Stephens RS, Myers G, Eppinger M, Bavoil PM. Divergence without difference: phylogenetics and taxonomy of *Chlamydia* resolved. *FEMS Immunol Med Microbiol.* 2009;55:115–9.
14. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol.* 2007;24:1596–9.