Antimicrobial Agents and Chemotherapy New Carbenicillin-Hydrolyzing β -Lactamase (CARB-7) from *Vibrio cholerae* Non-O1, Non-O139 Strains Encoded by the VCR Region of the *V. cholerae* Genome

Roberto Melano, Alejandro Petroni, Alicia Garutti, Héctor Alex Saka, Laura Mange, Fernando Pasterán, Melina Rapoport, Alicia Rossi and Marcelo Galas *Antimicrob. Agents Chemother.* 2002, 46(7):2162. DOI: 10.1128/AAC.46.7.2162-2168.2002.

	Updated information and services can be found at: http://aac.asm.org/content/46/7/2162
	These include:
REFERENCES	This article cites 23 articles, 15 of which can be accessed free at: http://aac.asm.org/content/46/7/2162#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: http://journals.asm.org/site/misc/reprints.xhtml To subscribe to to another ASM Journal go to: http://journals.asm.org/site/subscriptions/

Journals.ASM.org

New Carbenicillin-Hydrolyzing β-Lactamase (CARB-7) from *Vibrio cholerae* Non-O1, Non-O139 Strains Encoded by the VCR Region of the *V. cholerae* Genome†

Roberto Melano, Alejandro Petroni, Alicia Garutti, Héctor Alex Saka, Laura Mange, Fernando Pasterán, Melina Rapoport, Alicia Rossi,‡ and Marcelo Galas*

Servicio Antimicrobianos, Departamento de Bacteriología, Instituto Nacional de Enfermedades Infecciosas-ANLIS "Dr. Carlos G. Malbrán," Buenos Aires, Argentina

Received 24 August 2001/Returned for modification 12 January 2002/Accepted 7 April 2002

In a previous study, an analysis of 77 ampicillin-nonsusceptible (resistant plus intermediate categories) strains of *Vibrio cholerae* non-O1, non-O139, isolated from aquatic environment and diarrheal stool, showed that all of them produced a β -lactamase with a pI of 5.4. Hybridization or amplification by PCR with a probe for bla_{TEM} or primers for bla_{CARB} gene families was negative. In this work, an environmental ampicillin-resistant strain from this sample, ME11762, isolated from a waterway in the west region of Argentina, was studied. The nucleotide sequence of the structural gene of the β -lactamase was determined by bidirectional sequencing of a *Sau*3AI fragment belonging to this isolate. The gene encodes a new 288-amino-acid protein, designated CARB-7, that shares 88.5% homology with the CARB-6 enzyme; an overall 83.2% homology with PSE-4, PSE-1, CARB-3, and the *Proteus mirabilis* N29 enzymes; and 79% homology with CARB-4 enzyme. The gene for this β -lactamase could not be transferred to *Escherichia coli* by conjugation. The nucleotide sequence of the flanking regions of the *bla*_{CARB-7} gene showed the occurrence of three 123-bp *V. cholerae* repeated sequences, all of which were found outside the predicted open reading frame. The upstream fragment of the *bla*_{CARB-7} gene shared 93% identity with a locus situated inside *V. cholerae*'s chromosome 2. These results strongly suggest the chromosomal location of the *bla*_{CARB-7} gene, making this the first communication of a β -lactamase gene located on the VCR island of the *V. cholerae* genome.

Non-O1 Vibrio cholerae strains are natural inhabitants of aquatic environments. Although this group of microorganisms does not produce choleric diseases, a large cholera-like outbreak in Bangladesh and India was found to be caused by a V. cholerae non-O1 strain, which was named O139 because of its new O serogroup. Since 1981, ampicillin (AMP)-resistant V. cholerae strains have been sporadically isolated. These strains harbored plasmid-mediated β-lactamases, such as TEM-1 (8) and SAR-1 (19). More recently, a nonconjugative β -lactamase belonging to the CARB group, named CARB-6, was characterized from a clinical strain of V. cholerae non-O1, non-O139 (5). Among the carbenicillin-inactivating β -lactamases, the CARB family is the main group, including PSE-4 (CARB-1), PSE-1 (CARB-2), CARB-3, CARB-4, CARB-6, and N29, a β-lactamase from *Proteus mirabilis*. In addition, a new group of carbenicillinases, the RTG group, was proposed, which includes RTG-1 (P. mirabilis GN79 enzyme) and RTG-2 (CARB-5). Although these two chromosomal enzymes showed a low identity with members of the CARB family (44%), they may be considered to be the ancestors of this enzymatic group (6).

Some structural genes for CARB enzymes were located on resistance cassettes as part of integrons (18) that so far had been characterized as genetic elements capable of capturing and rearranging resistance genes (24). Integrons from class 1 were already observed in *V. cholerae* (7, 9). In addition, *V. cholerae* strains have a large chromosomal region, which acts as an integron-like structure, capturing genes from plasmids or via phages (15). This super integron is restricted to a region of about 10% of *V. cholerae* chromosome 2 (11) and is characterized by many 123- to 126-bp sequences of imperfect symmetry with a high of identity, named *V. cholerae* repeats (VCR). VCR flank one single gene or in some cases pair of genes (4). Most of the 216 open reading frames (ORFs) described in the VCR region have not shown homology with any known sequence, and only three appear to be involved in drug resistance (chloramphenicol [CMP] acetyltransferase, fosfomycin resistance protein, and glutathione transferase) (11).

In a previous work, we had performed susceptibility screening by disk diffusion methods of 669 isolates of *V. cholerae* non-O1, non-O139, which had been received at the National Institute of Infectious Diseases "Dr. C. G. Malbrán" between January 1993 and May 1998. Thirty percent of those isolates presented an AMP-nonsusceptible (resistant plus intermediate categories) phenotype. The analysis of a subset of 131 AMPnonsusceptible isolates showed that 57% of them produced a β -lactamase with a pI of 5.4 with high activity against aminoand carboxy-penicillins. For this group, the MICs at which 50 and 90% of isolates tested were inhibited (in micrograms per milliliter) were, respectively, as follows: AMP, 64 and 256, and ticarcillin, 128 and 256. Interestingly, the codifying gene for this β -lactamase did not belong to the TEM family. Moreover, no PCR products were obtained when specific primers for the

^{*} Corresponding author. Mailing address: Servicio Antimicrobianos, INEI-ANLIS "Dr. Carlos G. Malbrán," Av. Velez Sarsfield 563 (1281), Buenos Aires, Argentina. Phone and fax: 54-11-4303-2812. E-mail: mgalas@anlis.gov.ar.

[†] Dedicated to the memory of our dear Ali.

[‡] Deceased.

CARB-encoding genes were employed (L. Mange, H. Saka, A. Garutti, M. Galas, R. Melano, M. Rapoport, M. Caffer, and A. Rossi, 9th Int. Congr. Infect. Dis., abstr. 95.048, 2000).

In this work we analyzed an isolate belonging to that major group and we determined the nucleotide sequence of a new β -lactamase, named CARB-7, which in addition seems to be located on the VCR island.

MATERIALS AND METHODS

Bacterial strains and plasmids. *V. cholerae* non-O1, non-O139 (ME11762) was isolated from water samples from the Capilla waterway, Mendoza, Argentina. Pesc37 is an AMP-susceptible *V. cholerae* non-O1, non-O139 strain included in the initial sample of 669 isolates. Nalidixic acid-resistant *Escherichia coli* ER1793 was used as the recipient in conjugation experiments. *E. coli* DH5 α was used as the host for recombinant plasmids in cloning assays.

Plasmids pACYC184, encoding tetracycline and CMP resistance, and pBGS19, encoding kanamycin resistance (23), were used as cloning vectors.

Antimicrobial agents and susceptibility tests. The following antimicrobial agents were used and were obtained as standard laboratory powders from the sources indicated: AMP, Bagó; cefoxitin, Merck Sharp & Dohme; cephalothin, John Wyeth; piperacillin, Richmond; sulbactam, Pfizer; ticarcillin, Schering Plough; clavulanic acid, SmithKline Beecham; nalidixic acid, Winthrop.

MICs were determined by the agar dilution technique under *V. cholerae* NCCLS specifications (17), adopting the criteria for *Enterobacteriaceae* for β -lactam agents other than AMP.

Conjugative assays. Cells of both the donor and recipient strains were mixed on Luria-Bertani agar, and the mixture was incubated for 18 h at 25°C (3) or 35°C, as described previously (20). Mueller-Hinton agar supplemented with nalidixic acid (50 μ g/ml) and AMP (8 μ g/ml) was used to select the transconjugant cells.

DNA manipulations and PCR assays. Genomic DNA and plasmids were purified by employing the Wizard Genomic DNA Purification Kit and the Wizard Minipreps DNA Purification System (Promega, Madison, Wis.), respectively. The restriction enzymes as well as the DNA ligase were purchased from Gibco BRL (Gaithersburg, Md.). The genomic DNA was partially digested with *Sau*3AI and ligated with the pACYC184 vector linearized with *Bam*HI. The ligation mixture was electroporated into *E. coli* DH5 α competent cells using the Gene Pulser II Electroporation System (Bio-Rad) according to the manufacturer's recommendations. Selection of recombinant strains was achieved in Mueller-Hinton agar supplemented with AMP (16 µg/ml) and CMP (50 µg/ml).

Amplifications were performed with a Biometra thermal cycler in a final volume of 50 μ l containing 20 pmol of each primer (Gibco BRL), a 25 μ M concentration of each deoxynucleoside triphosphate, 1.5 mM MgCl₂, 2.5 U of *Taq* polymerase (Promega), and 1 μ l of DNA of ME11762 as the template. After a denaturation step of 10 min at 94°C, the amplifications were performed over 25 cycles, each one of which consisted of 30 s of denaturation at 94°C, 45 s of annealing at 60°C, and 30 s of extension at 72°C, with a final extension step of 5 min at 72°C. Seminested PCR was performed under the same conditions, using forward PD (5'-CTCTGATGCATGCCTTGCTACT-3') and reverse P3+ (5'-G CCTTATCAACGCGACTGTGAT-3') primers in the first round of amplification and forward PD and reverse P2+ (5'-GGCGTTGTCGTATCCCTCAAA T-3') primers in the second round. One microliter of amplimers obtained in the first round was employed as the template for the second round.

DNA sequencing and sequence analysis. DNA sequencing was performed on both strands by the method of Sanger et al. (22), using the BigDye terminator methodology (Applied Biosystem/Perkin-Elmer, Foster City, Calif.) with universal (M13/pUC forward and reverse 23-base sequencing primers) and sequencebased primers which, besides P2+, P3+, and PD, consisted of the following: P1, 5'-GGGCAACCCGTATTGAGATCACA-3'; P2-, 5'-GCCTGCATTGGTCA GTTCATAGGT-3'; P3-, 5'-CCGAGGCTTCAATGGCAGAA-3'; and P4, 5'-TGCGCGAAGTAGTGTTGTTG-3'. The sequences were analyzed in an ABI Prism 377 DNA sequencer (Applied Biosystem/Perkin-Elmer). Multiple nucleotide and protein sequence alignments were performed with the CLUSTAL facilities of the PCGENE software package (Intelligenetics, Inc.).

Nucleotide sequence accession number. The CARB-7 β -lactamase sequence has been submitted to GenBank. Its accession number is AF409092

RESULTS

Antibiotic susceptibility. Susceptibility testing of the ME11762 and Pesc37 isolates, respectively, determined the MICs (in micrograms per milliliter) of the following antibiotics: AMP, 256 and 0.5; AMP-sulbactam, 16 and 1; ticarcillin, 512 and 0.25; ticarcillin-clavulanic acid, 8 and 0.25; cephalothin, 2 and 0.25; piperacillin, 32 and 0.25; and cefoxitin, 8 and 1.

Conjugational transfer of resistance. To analyze the mobility of the AMP resistance determinant, the ME11762 isolate was mated with nalidixic acid-resistant *E. coli* ER1793. After several attempts, including assays performed at 25 or 37°C, no transconjugants were obtained.

Cloning and sequencing of bla_{CARB-7} gene. A Sau3AI-based library of the ME11762 isolate was prepared as indicated in Materials and Methods. *E. coli* DH5 α electrotransformants harboring recombinant plasmids were screened on selective agar containing AMP plus CMP. Plasmids from several AMP-and CMP-resistant colonies were analyzed, and the one with the smallest insert (approximately 2.2 kb), named pGMPS1, was subjected to restriction analysis (Fig. 1A).

To sequence the gene encoding AMP resistance, a SalI-EcoRV fragment from pGMPS1 (about 2.7 kb, including 0.5 kb of the pACyC184 vector) was directionally subcloned into the SalI-SmaI sites of pBGS19, rendering plasmid pGMPS2 (Fig. 1B). From this recombinant plasmid, three SacI-HindIII fragments were subsequently obtained and sequenced (Fig. 1B). To overlap the inner ends of these fragments, a PCR product obtained with primers P2+ and P2- was also sequenced. In addition, the sequencing of this fragment directly amplified from the chromosome allowed us to discard putative rearrangements of the Sau3AI fragments produced in the construction of the DNA library. The cloned DNA region of ME11762 strain resulted in a sequence of 2,214 nucleotides containing a unique ORF of 867 nucleotides (Fig. 1C and 2). Neither putative ribosome-binding site nor promoter regions were found. The comparison of this ORF with GenBank sequences showed a high nucleotide identity (88.6%) with the CARB-6 gene and an overall 82.5% identity with other genes belonging to the CARB family. This ORF encodes a deduced protein of 288 amino acids which showed 88.5% identity with CARB-6 (differing in 31 amino acid positions) and an overall identity of 83.4% with other enzymes of the CARB family (Fig. 3 and Table 1). The new enzyme, named CARB-7, contains the common motifs found in serine β-lactamases: the active site of the enzyme contained a Ser-Thr-Phe-Lys tetrad at positions 70 to 73 according to the standard numbering scheme of Ambler et al. (1), all seven conserved amino acid boxes (12), and specific conserved residues (Fig. 3). Alignment of amino acid sequences of 28 class A enzymes belonging to different β-lactamase-groups (TEM, SHV, CARB, PER, and CTX-M) showed a conserved Gly residue at position 144 (data not shown). Interestingly, an Asp residue at this position was found in the deduced amino acid sequence of CARB-7. Therefore, to confirm this change we designed a specific sequence-based primer, PD, and carried out a seminested PCR assay using PD and P2+ primers in the second round of amplification. The sequence of both strands (191 bp) of the PD-P2+ PCR frag-



FIG. 1. Strategy used for sequencing the bla_{CARB-7} gene and the DNA flanking regions. (A) Restriction map of the *Sau*3AI fragment from the ME11762 strain (black box). Only the ending *Sau*3AI sites are indicated. Grey boxes indicate DNA of pACYC184 vector. (B) The insert of pGMPS1 was liberated by digestion with *Sal*I and *Eco*RV and was subcloned into *SmaI-Sal*I sites of pBGS19 vector (white boxes), rendering pGMPS2; grey boxes represent approximately 500 bp that belong to pACYC184 vector. Inserts of plasmids derived of pGMPS2 were sequenced. (C) bla_{CARB-7} gene and its flanking DNA regions. The sense, length, and primers used for each sequence reaction are indicated by the black arrows. The grey box represents the bla_{CARB-7} gene, the white arrow represent its translational orientation, the flanking fragments are depicted as hatched boxes, and VCR are depicted as open boxes. Restriction sites used for subcloning into pBGS19 vector are indicated. Pf and Pr, universal primers (forward and reverse, respectively).

ment confirmed the presence of a GAT codon coding for an Asp residue at position 144 of the CARB-7 enzyme.

Analysis of flanking regions of the bla_{CARB-7} gene. The analysis of the nucleotide sequence of the flanking regions of the bla_{CARB-7} gene (887 nucleotides upstream and 460 nucleotides downstream) showed the occurrence of three 123-bp repeat sequences, two of them located on the upstream fragment and the other on the downstream fragment (Fig. 1C and 2). The three repeated sequences share an overall 97% identity with the reported consensus sequence of the VCR from *V. cholerae* (Fig. 4). As expected from this high degree of identity, the predicted secondary structure for these three repeated sequences showed stem-loop conformations having free-energy stability values in the range of -62.6 to -57.8 kcal/mol, which clearly resembles that from the previously published VCR consensus sequence (-58.6 kcal/mol) (Fig. 4) (4). The three VCR

are in the same orientation relative to one another, and all of them are found outside of the predicted ORF. Both flanking regions of the bla_{CARB-7} gene were compared with GenBank sequences. As is indicated in Fig. 2, 762 nucleotides belonging to the upstream fragment share 93% identity with locus VCA0424 (The Institute for Genomic Research), which is included within section 33 of *V. cholerae* chromosome 2 (Gen-Bank AE004376, nucleotides 7944 to 8706). The downstream fragment mainly showed high homology with other VCR sequences.

DISCUSSION

Both *V. cholerae* O1 and non-O1 are naturally susceptible in vitro to broad-spectrum penicillins and cephalosporins. However, strains producing β -lactamases such as SAR-1 (pI 4.9),

gato atca gego acao gtta acao	cttta agcad ctgta ggcto atcgt gagco aaaco	atgg ggtt aaat gaaa tgcco cagt gcct	aaaca atgao gtgaa totga gtoto catgo totaa caaga	aaatg gccgo atgto agtta gtcgo cgtgo attoa	gggat cggt cggt attg gcaat gcaat gtga aagga actgt	ttta ggtt ccti ccto gtga aaach	ttgcagi gcagi ttggo actao atati gttta cgcgi	aatto gatto ggtto tgtto atcgo agtto tagco	gaaca cgtgt ccgtt ccgtt ctga cgga cgga ggga g	aaggi cttg ccage agte geta geta caate ccage	ctca agtti gtti cgcgo cota ctti gatti	caaa cggt cgtc cgag atga acat cag attg	tagca ggtai ggtai gttai aacgo tggco ttggi agcco	acgca attca gtgtt gcaat ctagt ctato gcgat	aca <u>ta</u> aata tgtto tgtg tgtg catto tggt	aacaa cagco ttgt ggcto tetgi taaai gatti tacgo	aacgo cccti ttgto caagi tcgci tgaa tgati gttgi	cetea cgcto cgcto cgcto cgtto cgtaa ageet caaa ttgto	aagao goota ittga agtti igcat atgta	gggad cgtta aatga agcaq ccagt ccagt ccagt aaagt gagtt	atgc agc gttt gttg cagt cagt tgtt tgtt	saaco staat ggcag ggtgc sttgc tcat <u>gaaqc</u> tgtta	gogto cocat gtgta cocttt coctoa caggt cttaa	igagt aaaa atctg actt attga aaaa agca <u>t</u> gttgt	100 200 300 400 500 600 700 800
cago	eccet	ttag	gegg	gcgtt	tage	cttai	tttca	agtgo	cctaa	atge	tttaa	agtti	aagat	cgtat	tgeea	aatca	aatat	ttato	idddi	caca	aac	M ATG	K AAG	S TCT	3 896
L	L	v	F	A	L	L	М	P	S	v	v	F	A	S	S	S	K	F	Q	S	v	Е	Q	E	28
TTG	TTG	GTA	TTT	GCG	CTT	TTA	ATG	CCA	TCT	GTA	GTT	TTT	GCA	AGC	AGT	TCA	AAA	TTT	CAA	TCA	GTT	GAA	CAA	GAA	971
I ATT	K AAG	G GGA	I ATT	E GAG	S TCT	S TCA	L CTC	S TCT	A GCT	R CGT	I ATA	G GGA	V GTC	A GCC	I ATT	L TTG	D GAT	T ACT	Q CAA	N AAT	G GGC	E GAA	S AGC	W TGG	53 1046
D	Y	N	G	D	Q	R	F	P	L	т	S	Т	F	К	т	I	A	С	A	K	L	L	Y	D	78
GAT	TAT	AAT	GGT	GAT	CAA	CGA	TTT	CCA	TTA	ACA	AGT	ACT	TTC	AAA	ACA	ATA	GCT	TGT	GCT	AAG	TTG	CTG	TAT	GAT	1121
A GCA	E GAG	H CAT	G GGG	K AAA	V GTT	N AAT	L CTC	N AAT	S AGT	T ACA	V GTT	E GAG	I ATT	K AAG	K AAA	A GCA	D GAT	L CTT	V GTT	T ACG	Y TAT	S TCG	P CCT	V GTA	103 1196
L	E	K	Q	V	G	K	P	I	T	L	S	D	A	C	L	A	T	M	T	T	S	D	N	Т	128
TTA	GAA	AAG	CAA	GTA	GGT	AAA	CCA	ATA	ACG	CTC	TCT	GAT	GCA	TGC	UTT	GUT	ACT	ATG	AUA	ACA	AGC	GAC	AAT	ACA	12/1
A GCA	A GCC	N AAT	I ATT	V GTT	I ATA	N AAT	A GCT	V GTC	G GGT	D GAT	P CCT	K AAA	S AGC	I ATT	T ACT	D GAT	F TTT	L CTG	R AGA	Q CAA	I ATT	G GGT	D GAC	K AAA	153 1346
E	Т	R	L	D	R	V	E	P	Е	L	Ν	E	G	K	L	G	D	L	R	D	Т	Т	Т	Р	178
GAA	ACT	CGT	CTA	GAT	CGT	GTC	GAG	CCT	GAG	CTC	AAT	GAA	GGT	AAA	CTC	GGT	GAT	TTG	AGG	GAT	ACG	ACA	ACG	CCT	1421
N	А	Ι	Т	S	Т	L	Ν	Q	L	L	F	G	S	Т	L	S	Е	A	S	Q	Κ	К	L	E	203
AAT	GCA	ATA	ACC	AGC	ACG	TTA	AAT	CAA	TTA	TTA	TTT	GGT	TCC	ACA	TTA	TCT	GAA	GCT	AGT	CAG	AAA	AAA	TTA	GAG	1496
S TCT	W TGG	M ATG	V GTG	N AAC	N AAT	Q CAA	V GTT	T ACG	G GGT	N AAT	L TTA	L TTG	R AGG	S TCA	V GTA	L TTG	P CCA	V GTG	K AAG	W TGG	S AGT	I ATT	A GCT	D GAT	228 1571
R CGC	S TCA	G GGA	A GCA	G GGT	G GGA	F TTT	G GGT	A GCT	R AGG	S AGT	I ATT	T ACA	A GCG	I ATT	V GTG	W TGG	S AGT	E GAA	e gaa	K AAA	K AAA	P CCG	I ATT	I ATC	253 1646
V	S	I	Y	L	A	Q	Т	E	А	S	Μ	A	Е	R	N	D	A	I	v	K	I	G	R	S	278
GTA	AGT	ATT	TAT	CTA	GCT	CAA	ACC	GAG	GCT	TCA	ATG	GCA	GAA	CGA	AAT	GAT	GCG	ATA	GTT	AAG	ATT	GGT	CGT	TCA	1721
I ATT	F TTT	E GAA	V GTT	Y TAT	T ACA	S TCA	Q CAG	S TCG	R CGT	- TGA	taa	ggc <u>t</u>	aacaa	acgo	cetea	aaga	ggga	ctgto	caaco	leded	ggogt	ttec	cagto	cct	288 1809
$\frac{\texttt{atgagccgcggtggttactgttgttgttgtgtttaggtttagtgttttgcgttgtcagccccttaggcgggggttagcttcttgaagcgaaaattatcaaaa 19}{\texttt{tataagccattgcttgacttgacttgacttgacttggtcagcaccgattggttcagctgttgtgtgtagaagtatgaaag 20} \\ \texttt{tcattagttcaaagccgctgaaaacgagcaagcctcgatgctcagcatcgttggatgtcaacactcaaagcctggtttcacaaaaggctgcacta 22} \\ cgctgtgactcaatacggttgccctcgatcactatcctattttgccgtggtccccgatattcctcgctggggatcactcac$										1909 2009 2109 2209 2214															

FIG. 2. Nucleotide sequence of the 2,214-bp fragment of pGMPS2 containing the CARB-7 β -lactamase coding region. The deduced amino acid sequence of the ORF proposed to encode CARB-7 β -lactamase is represented in one-letter code. VCR on the flanking sequences of bla_{CARB-7} gene are underlined. The fragment showing high homology with the *V. cholerae* chromosome is indicated by the grey box (GenBank accession number AE004376 [nucleotides 7944 to 8706]). Restriction sites used for subcloning of bla_{CARB-7} gene are double underlined.

TEM-1 (pI 5.4), or CARB-6 (pI 5.35), which conferred resistance to amino- and carboxy-penicillins, were previously described (5, 8, 19). We have reported here the sequence of a new β -lactamase gene isolated from a *V. cholerae* non-O1, non-O139 strain, which shares an overall 85.5% identity with the genes coding for several CARB enzymes. Moreover, from all the sequences currently comprised in the GenBank, only the CARB genes showed a relevant homology with the encoding gene for the new β -lactamase. In addition, the susceptibility profile conferred by the new β -lactamase suggests a strong activity against amino- and carboxy-penicillins, which resembled the phenotype conferred by PSE-1 (CARB-2) or PSE-4 (CARB-1) β -lactamases in *Pseudomonas aeruginosa* (14). Therefore, we propose that the new enzyme is a member of the CARB group and name it CARB-7.

Despite the fact that both the nucleotide and deduced amino acid sequences for CARB-7 showed high similarity with the those of the CARB enzymes, two relevant differences must be pointed out. First, as could be expected from having the lowest overall similarity percentage observed with other CARB genes, there is a poor homology between the bla_{CARB-7} gene and the reverse specific primer (OC-2) reported by Arlet and Philippon (2) for the identification of members of the bla_{CARB} gene family. This fact can explain the lack of amplification in the PCR assays carried out in our preliminary work for the molecular identification of β-lactamases (Mange et al., 9th ICID). It points out the need to design new primers that can detect this new member of the CARB family. Second, the CARB group includes enzymes belonging to the molecular class A β-lactamases, for which several conserved residues or motifs have been identified. Comparing these conserved motifs on the deduced sequence for CARB-7 with those of B-lactamases from the major families classified into class A, such as TEM, SHV, CARB, CTX-M, and PER enzymes, we observed a change, Gly to Asp, at position 144 (Ambler classification scheme) of the CARB-7 enzyme. This is, to our knowledge, the first report of this amino acidic variant. It would be located far from the active site, as can be inferred from the crystallo-

		Box-I	
PSE4	MKFLLAFSLLIPSVVFA	SSSKFOOVEODVKAIEVSLSARIGVSVLDTONGEYWDY	60
PSE1	MKFLLAFSLLIPSVVFA	SSSKFQQVEQDVKAIEVSLSARIGVSVLDTQNGEYWDY	60
CARB3	MKFLLAFSLLIPSVVFA	SSSKFQQVEQDVKAIEVSLSARIGVSVLDTQNGEYWDY	60
N29	MKFLLVFSLLIPSVVFA	SSSKFRQVEQDVKAIEVSLSARIGVSVLDTQNGEYWDY	60
CARB4	MKLLLVFSLLIPSMVFA	NSSKFQQVEQDAKVIEASLSAHIGISVLDTQTGEYWDY	60
CARB6	MKFLLAFSLLIPSVVFA	SSSKFQQVEQDVKAIEVSLSARIGVSVLDTQNGEYWDY	60
CARB7	MKSLLVFALLMPSVVFA	SSSKFQSVEQEIKGIESSLSARIGVAILDTQNGESWDY	60
	** **.*.**.**.	.****. ***. * ** ***.******.** ***	
	Box-II	Box-III Box-IV	
PSE4	NGNQRFPLTSTFKTIAC	AKLLYDAEQGKVNPNSTVEIKKADLVTYSPVIEKQVGQ	115
PSE1	NGNQRFPLTSTFKTIAC	AKLLYDAEQGKVNPNSTVEIKKADLVTYSPVIEKQVGQ	115
CARB3	NGNQRFPLTSTFKTIAC	AKLLYDAEQGKVNPNSTVEIKKADLVTYSPVIEKQVGQ	115
N29	NGNQRFPLTSTFKTIAC	AKLLYDAEQGKVNPNSTVEIKKADLVTYSPVIEKQVGQ	115
CARB4	NGNQRFPLTSTFKTIAC	AKLLYDAEQGEINPKSTIEIKKADLVTYSPVIEKQVGQ	115
CARB6	NGNQRFPLTSTFKTIAC	AKLLYDAEQGKVNPNSTVEIKKADLVTYSPVIEKQVGQ	115
CARB7	NGDQRFPLTSTFKTIAC	AKLLYDAEHGKVNLNSTVEIKKADLVTYSPVLEKQVGK	115
	** . * * * * * * * * * * * * * * *	*******.** .**.**********************	
		Box-V	
PSE4	AITLDDACFATMTTSDN	TAANIILSAVGGPKGVTDFLRQIGDKETRLDRIEPDLN	170
PSE1	AITLDDACFATMTTSDN	TAANIILSAVGGPKGVTDFLRQIGDKETRLDRIEPDLN	170
CARB3	AITLDDACFATMTTSDN	TAANIILSAVGGPKGVTDFLRQIGDKETRLDRIEPDLN	170
N29	AITLDDACFATMTTSDN	TAANIILSAVGGSKGVTDFLRQIGDKETRLDRIEPDLN	170
CARB4	AITLDDACFATMTTSDN	AAANIILNALGGPESVTDFLRQIGDKETRLDRIEPELN	170
CARB6	AITLDDACFATMTTSDN	TAANIILSAVGGPKGVTDFLRQIGDKETRLDRIEPDLN	170
CARB7	PITLSDACLATMTTSDN	TAANIVINAVGDPKSITDFLRQIGDKETRLDRVEPELN	170
	*** *** ******	· * * * * · · · * · * 1 · · · · * * * * * * * * * * * * * * *	
		Box-VI	
PSE4	EGKLGDLRDTTTPKAIA	STLNKFLFGSALSEMNQKKLESWMVNNQVTGNLLRSVL	225
PSE1	EGKLGDLRDTTTPKAIA	STLNKFLFGSALSEMNQKKLESWMVNNQVTGNLLRSVL	225
CARB3	EGKLGDLRDTTTPKAIA	STLNKLLFGSALSEMNQKKLESWMVNNQVTGNLLRSVL	225
N29	EGKLGDLRDTTTPKAIA	STLNKFLFGSALSEMNKKKLESWMVNNQVTGNLLRSVL	225
CARB4	EGKLGDLRDTTTPNAIV	NTLNELLFGSTLSQDGQKKLEYWMVNNQVTGNLLRSVL	225
CARB6	EGKLGDLRDTTTPKAIA	STLNQLLFGSTLSEASQKKLESWMVNNQVTGNLLRSVL	225
CARB7	EGKLGDLRDTTTPNAIT	STLNQLLFGSTLSEASQKKLESWMVNNQVTGNLLRSVL	225
	Box-VII		
PSE4	PAGWNIADRSGAGGFGA	RSITAVVWSEHQAPIIVSIYLAQTQASMEERNDAIVKI	280
PSEI	PAGWNIADRSGAGGFGA	RSITAVVWSEHQAPIIVSIYLAQTQASMAERNDAIVKI	280
CARB3	PAGWNIADRSGAGGFGA	RSITAVVWSEHQAPIIVSIYLAQTQASMAERNDAIVKI	280
NZ9	PAGWNIADRSGAGGEGA	RSITAVVWSEHQAPIIVSIYLAQTQASMAERNDAIVKI	280
CARB4	PEGWNIADRSGAGGFGA	RSITAVVWSEAQSPIIVSIILAQTEASIADRNDAIVKI	280
CARBO	PVRWSTADRSGAGGEGA	RSITALVWSEERRIIIVSIILAQTEASMAERNDAIVKI	280
CARD /	* * * **********	*****.****************************	200
PSE4	GHSIFDVYTSQSR 2	93	
PSE1	GHSIFDVYTSQSR 2	93	
CARB3	GHSIFDVYTSQSR 2	93	
N29	GRSIFDVYTSQSR 2	93	
CARB4	GRSIFEVYSSQSR 2	93	
CARB5	GREEFEVITEQSR 2	93	
CHIG8 /	* *** ** ****	33	

FIG. 3. Alignment of the deduced amino acid sequence of CARB-7 with those of the closely related class A β -lactamases PSE-4 (CARB-1), PSE-1 (CARB-2), CARB-3, CARB-4, *P. mirabilis* N29, and CARB-6. The shadowed boxes (I to VII) correspond to amino acid boxes conserved in all penicillin-recognizing enzymes (12). The arrow indicates amino acid position 139, with the change G \rightarrow D. Asterisks indicate identical residues. Dots indicate conserved amino acids. Sequences are numbered as described by Ambler et al. (1).

graphic structure reported for the TEM-1 β -lactamase (25). The strong activity against amino- and carboxy-penicillins displayed by CARB-7 appears to support the assumption that Asp144 is not involved in the enzymatic activity. Strikingly, the two Gly residues at positions 143 and 144 are highly conserved among the Ambler class A enzymes. Therefore, this nonconservative change of a nonpolar residue to an acidic one at position 144 of CARB-7 could imply a conformational change in the secondary structure of the enzyme, which could additionally have evolutionary significance.

Although initially the CARB enzymes were assumed to be confined to *P. aeruginosa* strains (10), some of these β -lactamases were isolated from enterobacteria (16), Alcaligenes xylosoxidans (CARB-2) (13), and from an isolate of V. cholerae non-O1, non-O139 (CARB-6) (5). In this study, CARB-7 was isolated from a V. cholerae non-O1, non-O139 strain from a sample of 77 AMP-resistant isolates that were producers of a β-lactamase with a pI of 5.4. The bla_{CABB-7} gene was not transferred to the E. coli ER1793 strain, regardless of the fact that several attempts employing different mating temperatures were made. This fact was previously reported for others CARB enzymes (5) and could be interpreted as either a location of the CARB gene on a nonconjugative plasmid or on the chromosome. This last possibility is strongly supported by two facts. First, three 123-bp repeats were detected on the flanking sequence of the bla_{CABB-7} gene showing a 97% overall similarity with the consensus sequence for VCR (4), which are restricted to a region of about 10% of V. cholerae's chromosome 2 (11). Second, the upstream fragment of the bla_{CARB-7} gene shared 93% identity with locus VCA0424 (The Institute for Genomic Research), situated inside section 33 of V. cholerae's chromosome 2. The assumption of a chromosomal location for the bla_{CARB-7} gene supports the notion that the VCR island functions as a "mega-integron." To date, only in vitro works have demonstrated that intl4 V. cholerae integrase is active for sitespecific cassette recombination (21), without in vivo proof of gene capture. Therefore, this would be the first evidence of in vivo capture of a gene by this system.

All the VCR-associated genes described to date are transcribed from promoter sequences located immediately upstream of the ORFs (C. Clark, L. Purins, P. Kaewrakon, and P. Manning, Letter, Mol. Microbiol. **26**:1137-1138, 1997). However, putative ribosome-binding site or promoter regions (regarding *E. coli* consensus sequences) were not found in the upstream region of the bla_{CARB-7} gene. These facts suggest

ΓABLE 1. Amino acid identity of CARB-7 with several related class A β -lact

0 I .	% Identity with:										
β-Lactamase	PSE-4 ^a	$PSE-1^b$	CARB-3	N29	CARB-6	CARB-4					
PSE-4											
PSE-1	99.7										
CARB-3	99.3	99.7									
N29	97.9	98.2	97.9								
CARB-6	93.8	94.1	94.4	93.1							
CARB-4	85.1	85.1	85.4	84.7	85.1						
CARB-7	83.0	83.3	83.7	83.0	88.5	78.8					

^a Also designated CARB-1.

^b Also designated CARB-2.

Α		
VCR1	TAACAAACGCCTCAAGAGGGACTGTCAACGCGTGGAGTTTCCAGTCCCAATGAGCCGCGGTG	GTT 65
VCR2	TAACAAACGCCTCAAGAGGGACTGTCAACGCGTAGCGTTTCCAGTCCCATTGAGCCGCGATG	GTT 65
VCR3	TAACAAACGCCTCAAGAGGGACTGTCAACGCGCGGCGTTTCCAGTCCCTATGAGCCGCGGTG	GTT 65
Consensus	TAACAAACGCCTCAAGAGGGACTGTCAACGCGTGGCGTTTCCAGTCCCATTGAGCCGCGGTG	GTT 65
VCP1		123
VCR2	ACCOTTCTTCTCTTTCACTTTCCTCTTA-TCCCTTCTCACCCCCTTACCCCCCCTTACCCCCCCTTACCCCCC	23
VCR3	ACTGTTGTTGTGTTTTGAGTTTTAGTGTTT-TGCGTTGTCAGCCCCTTAGGCGGGCGTTAG	123
Consensus	TCGGTTGTTGTGTTTGAGTTTAGTGGTAATGCGTTGCCAGCCCCTTAGGCGGCGTTAG 1 * **********************************	.24





FIG. 4. Nucleotide sequence of the 123-bp element. (A) Multiple alignment of nucleotide sequence of the three VCR found on the flanking regions of the bla_{CARB-7} gene with a proposed VCR consensus sequence (4). Identical bases are indicated by an asterisk. The alignment was generated with the multiple-alignment program CLUSTAL. (B) Proposed secondary structure for each VCR sequence, determined with DNASIS software (version 2.5).

2168 MELANO ET AL.

that the transcription may occur through a promoter located more distantly of the bla_{CARB-7} gene.

Most of the 216 ORFs situated on the VCR island have no homology to other sequences, although three genes were identified that may be involved in resistance to drugs other than β -lactam antibiotics (11). Therefore, this is the first report of a β -lactamase within the VCR region.

ACKNOWLEDGMENTS

We thank Mariana Miranda for her assistance in technical work. We thank N. Binsztein for providing the *V. cholerae* non-O1, non-O139 collection and for reading the manuscript and thank M. I. Caffer for serotyping of the strains.

REFERENCES

- Ambler, R. P., A. F. W. Coulson, J. M. Frere, J. M. Ghuysen, B. Joris, M. Forsman, R. C. Levesque, G. Tirabi, and S. G. Waley. 1991. A standard numbering scheme for the class A β-lactamases. Biochem. J. 276:269–272.
- Arlet, G., and A. Philippon. 1989. Construction by polymerase chain reaction and intragenic DNA probes for three main types of transferable β-lactamases (TEM, SHV, CARB). FEMS Microbiol. Lett. 82:19–26.
- Barja, J. L., Y. Santos, I. Huq, R. Colwell, and A. E. Toranzo. 1990. Plasmids and factors associated with virulence in environmental isolates of *Vibrio cholerae* non-O1 in Bangladesh. J. Med. Microbiol. 33:107–114.
- Barker, A., C. Clark, and P. Manning. 1994. Identification of VCR, a repeated sequence associated with a locus encoding a hemagglutinin in *Vibrio cholerae* O1. J. Bacteriol. 176:5450–5458.
- Choury, D., G. Aubert, M. Szajnert, K. Azibi, M. Delpech, and G. Paul. 1999. Characterization and nucleotide sequence of CARB-6, a new carbenicillinhydrolyzing β-lactamase from *Vibrio cholerae*. Antimicrob. Agents Chemother. 43:297–301.
- Choury, D., M.-F. Szajnert, M.-L. Joly-Gillou, K. Azibi, M. Delpech, and G. Paul. 2000. Nucleotide sequence of the bla_{RTG-2} (CARB-5) gene and phylogeny of a new group of carbenicillinases. Antimicrob. Agents Chemother. 44:1070–1074.
- Dalsgaard, A., A. Forslund, O. Serichantalergs, and D. Sandvang. 2000. Distribution and content of class 1 integrons in different *Vibrio cholerae* O-serotype strains isolated in Thailand. Antimicrob. Agents Chemother. 44:1315–1321.
- Dupont, M., M. Jouvenot, G. Couetdic, and Y. Michel-Briand. 1985. Development of plasmid-mediated resistance in *Vibrio cholerae* during treatment with trimethoprim-sulfamethoxazole. Antimicrob. Agents Chemoter. 27:280–281.
- Falbo, V., A. Carattoli, F. Tosini, C. Pezzella, A. Dionisi, and I. Luzzi. 1999. Antibiotic resistance conferred by a conjugative plasmid and a class I integron in *Vibrio cholerae* O1 El Tor strains isolated in Albania and Italy. Antimicrob. Agents Chemother. 43:693–696.
- 10. Hedges, R., and M. Matthew. 1979. Acquisition by Escherichia coli of plas-

mid-borne β -lactamases normally confined to *Pseudomonas* spp. Plasmid **2:**269–278.

- 11. Heidelberg, J., J. Eisen, W. Nelson, R. Clayton, M. Gwinn, R. Dodson, D. Haft, E. Hickey, J. Peterson, L. Umayam, S. Gill, K. Nelson, T. Read, H. Tettelin, D. Richardson, M. Ermolaeva, J. Vamathevan, S. Bass, H. Qin, I. Dragoi, P. Sellers, L. McDonald, T. Utterback, R. Fleishmann, W. Nierman, O. White, S. Salzberg, H. Smith, R. Colwell, J. Mekalanos, J. Venter, and C. Fraser. 2000. DNA sequence of both chromosomes of the cholera pathogen Vibrio cholerae. Nature 406:477–483.
- Joris, B., J. Ghuysen, G. Dive, A. Renard, O. Dideberg, P. Charlier, J. Frère, J. Kelly, J. Boyington, P. Moews, and J. Knox. 1988. The active-site-serine penicillin-recognizing enzymes as members of the *Streptomyces* R61 DDpeptidase family. Biochem. J. 250:313–324.
- Labia, R., M. Guionie, M. Barthélémy, and A. Philippon. 1981. Properties of three carbenicillin-hydrolyzing β-lactamases (CARB) from *Pseudomonas* aeruginosa: identification of a new enzyme. J. Antimicrob. Chemother. 7:49– 56.
- Livermore, D. M. 1995. β-Lactamases in laboratory and clinical resistance. Clin. Microbiol. Rev. 8:557–584.
- Mazel, D., B. Dychinco, V. Webb, and J. Davies. 1998. A distinctive class of integron in the Vibrio cholerae genome. Science 280:605–608.
- Medeiros, A., R. Hedges, and G. Jacoby. 1982. Spread of a "Pseudomonasspecific"β-lactamase to plasmids of enterobacteria. J. Bacteriol. 149:700– 707.
- National Committee for Clinical Laboratory Standards. 2000. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically, 5th ed., vol. 20, no. 2. Approved standard M7-A5. National Committee for Clinical Laboratory Standards, Wayne, Pa.
- Recchia, G., and R. Hall. 1995. Gene cassettes: a new class of mobile element. Microbiology 141:3015–3027.
- Reid, A., and S. Amyes. 1986. Plasmid penicillin resistance in Vibrio cholerae: identification of new β-lactamase SAR-1. Antimicrob. Agents Chemother. 30:245–247.
- Rice, L. B., and R. A. Bonomo. 1996. Genetic and biochemical mechanisms of bacterial resistance to antimicrobial agents, p. 453–501. *In* M. Lorian (ed.), Antibiotics in laboratory medicine, 4th ed. Williams & Wilkins, Baltimore, Md.
- Rowe-Magnus, D., A. Guerout, P. Ploncard, B. Dychinco, J. Davies, and D. Mazel. 2001. The evolutionary history of chromosomal super-integrons provides an ancestry for multiresistant integrons. Proc. Natl. Acad. Sci. 98:652– 657.
- Sanger, S., S. Micklen, and A. Coulson. 1977. DNA sequencing with chainterminating inhibitors. Proc. Natl. Acad. Sci. USA 74:5463–5467.
- Spratt, B. G., P. J. Hedge, S. te Heesch, A. Edelman, and J. K. Broome-Smith. 1986. Kanamycin-resistant vectors that are analogues of plasmids pUC8, pUC9, pEMBL8 and pEMBL9. Gene 41:337–341.
- Stokes, H. W., and R. M. Hall. 1989. A novel family of potentially mobile DNA elements encoding site-specific gene-integration functions: integrons. Mol. Microbiol. 3:1669–1683.
- Strynadka, N. C. J., H. Adachi, S. E. Jensen, K. Johns, A. Sielecki, C. Betzel, K. Sutoh, and M. N. G. James. 1992. Molecular structure of the acyl-enzyme intermediate in β-lactam hydrolysis at 1.7 Å resolution. Nature (London) 359:700–705.